

Phosphate desorption from the surface of soil mineral particles by a phosphate-solubilizing fungus

Nelson Walter Osorio & Mitiku Habte

Biology and Fertility of Soils
Cooperating Journal of International
Society of Soil Science


ISSN 0178-2762

Volume 49

Number 4

Biol Fertil Soils (2013) 49:481-486

DOI 10.1007/s00374-012-0763-5



Biology and Fertility of Soils

Volume 49 • Number 4 • May 2013

ISSN-AISS-IBG - Cooperating Journal of the
International Society of Soil Science

REVIEW

Changes in soil microbial properties with no-tillage in Chinese cropping systems
M. Huang · L. Jiang · Y. Zou · S. Xu · G. Deng 373

ORIGINAL PAPERS

Salinity reduces the ability of soil microbes to utilize cellulose
B. Elmajdoub · P. Marschner 379

Continuous manuring combined with chemical fertilizer affects soil microbial residues in a Mollisol
X. Ding · X. Han · X. Zhang · Y. Qiao · Y. Liang 387

Effect of organic-complexed superphosphates on microbial biomass and microbial activity of soil
C. Giovannini · J.M. Garcia-Mina · C. Ciavatta · C. Marzadori 395

Early changes due to sorghum biofuel cropping systems in soil microbial communities and metabolic functioning
J. Cotton · V. Acosta-Martinez · J. Moore-Kucera · G. Burov 403

Organic amendments differ in their effect on microbial biomass and activity and on P pools in alkaline soils
M.A. Malik · K.S. Khan · P. Marschner · S. Ali 415

Home-field advantage of litter decomposition and nitrogen release in forest ecosystems
Q. Wang · M. Zhong · T. He 427

Effects of novel bioorganic fertilizer produced by *Bacillus amyloquelaciens* W19 on antagonism of *Fusarium* wilt of banana
B. Wang · J. Yuan · J. Zhang · Z. Shen · M. Zhang · R. Li · Y. Ruan · Q. Shen 435

For continuation of table of contents, see inside back cover

Further articles can be found at www.springerlink.com

Indexed in Science Citation Index, Science Citation Index Expanded (SciSearch), Journal Citation Reports/Science Edition, SCOPUS, Chemical Abstracts Service (CAS), Google Scholar, EBSCO, CSA, ProQuest, CAB International, Academic OneFile, AGRICOLA, Biological Abstracts, BIOSIS, CAB Abstracts, Current Abstracts, Current Contents/Agriculture, Biology & Environmental Sciences, Elsevier Biobase, EMBiology, Gale, Geobase, GeoRef, Global Health, International Bibliography of Book Reviews (IBR), International Bibliography of Periodical Literature (IBZ), OCLC, SCImago, Summon by Serial Solutions, VINITI - Russian Academy of Science, Zoological Record

Instructions for Authors for *Biol Fertil Soils* are available at www.springer.com/00374

 Springer

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Phosphate desorption from the surface of soil mineral particles by a phosphate-solubilizing fungus

Nelson Walter Osorio · Mitiku Habte

Received: 7 September 2012 / Revised: 14 November 2012 / Accepted: 27 November 2012 / Published online: 15 December 2012
© Springer-Verlag Berlin Heidelberg 2012

Abstract High phosphate (Pi) sorption in soils is a serious limiting factor for plant productivity and Pi fertilization efficiency, particularly in highly weathered and volcanic ash soils. In these soils, the sorbed Pi is so strongly held on the surfaces of reactive minerals that it is not available for plant root uptake. The use of phosphate-solubilizing microorganisms (PSM) capable of Pi desorption seems to be a complementary alternative in the management of these soils. The aim of this study was to evaluate the effectiveness of the soil fungus *Mortierella* sp., a known PSM, to desorb Pi from four soil minerals differing in their Pi sorption capacity. The fungus was effective in desorbing Pi from all tested minerals except from allophane, and its desorption depended on the production of oxalic acid. The effectiveness of the fungus to desorb Pi was ranked as montmorillonite > kaolinite > goethite > allophane. The quantity of desorbed Pi increased by increasing the amount of sorbed Pi. The Pi sorption capacity expressed as $P_{0.2}$ value (amount of P required to increase a solution P concentration up to 0.2 mg L^{-1}) was a good indicator of the effectiveness of *Mortierella* sp. to desorb Pi from soil minerals.

Keywords *Mortierella* · Phosphorus · Montmorillonite · Kaolinite · Goethite · Allophane

Introduction

Phosphorus sorption is a process by which bioavailable phosphate (Pi) is removed from a soil solution and retained on the surface of soil minerals (Do Carmo Harta and Torrent 2007). Phosphate sorption is particularly strong on iron and aluminum hydrous oxides (e.g., goethite) that predominate in the highly weathered soils of the tropics (Jackman et al. 1997; Onweremadu 2007). In soils derived from volcanic materials, allophane, ferrihydrite, and goethite are responsible for the strong Pi sorption of these soils (Shoji et al. 1993; Jackman et al. 1997; Vistoso et al. 2012).

Many organic acids are effective in either reducing Pi sorption or desorbing Pi (Guppy et al. 2005), and, according to Hue (1991) and Bolan et al. (1994), their effectiveness in decreasing soil Pi sorption could be ranked as tricarboxylic > dicarboxylic > monocarboxylic acids. Organic acids such as citric, oxalic, gluconic, and malic, among others are produced by roots (Kirk et al. 1999; Radersma and Grierson 2004; Nwoke et al. 2008; Corrales et al. 2007; Li et al. 2009) and soil microorganisms (Gyaneshwar et al. 2002; Reddy et al. 2002; Welch et al. 2002; Chen et al. 2006; Pandey et al. 2006; Puente et al. 2009; Bashan et al. 2012). Therefore, soil microorganisms may desorb Pi from the surfaces of reactive particles of soils as reported by He and Zhu (1998); Hoberg et al. (2005), and He et al. (2002).

The phenomenon of Pi desorption by organic anions is widely accepted as a mechanism to enhance Pi availability in the rhizosphere, and this occurs onto soil surface by ligand exchange (Trove et al. 2003; Sato and Comerford 2006; Bashan et al. 2012). However, the concentration of these organic acid/anions is relatively low in most soils because they can be precipitated with free ions (e.g., Al^{3+} , Fe^{3+} , Ca^{2+}), sorbed on the surface of soil minerals, and/or used as carbon sources by soil microorganisms (Jones et al. 2003). Thus, competent rhizosphere microflora capable of

N. W. Osorio (✉)
Universidad Nacional de Colombia, Calle 59A No. 63-20,
Of. 14-216,
050034, Medellín, Colombia
e-mail: nwsorio@unal.edu.co

M. Habte
University of Hawaii, St. John 102, 3190 Maile Way,
96822, Honolulu, HI, USA
e-mail: mitiku@hawaii.edu

producing organic acids or anions may play an important role in the management of Pi deficiency in high Pi-sorbing soils. He et al. (2002) found that a bacterium identified as *Moraxella* sp. was capable of desorbing Pi from the surface of goethite, montmorillonite, and kaolinite.

The hypothesis tested was that the effectiveness of a phosphate-solubilizing microorganism (PSM) may desorb Pi from the surface of soil mineral particles, but such Pi desorption is controlled by the Pi sorption capacity (measured as the amount of P required to increase soluble P concentration up to 0.2 mg L⁻¹) and the extent of saturation of Pi sorption sites in these mineral particles. The objective of this study was to determine the effectiveness of the fungus *Mortierella* sp., a known phosphate-solubilizing fungus, in desorbing Pi from the surfaces of soil reactive particles (allophane, goethite, kaolinite, and montmorillonite) widely differing in their Pi sorption capacity.

Material and methods

Fungal inoculum

The fungus *Mortierella* sp. is a known effective PSM that was originally isolated from an Andisol of Hawaii (Osorio and Habte 2001) and has been cultured on yeast mannitol agar (YMA) medium and stored on YMA slants at 4 °C. For this study, the fungus was multiplied in Petri dishes on YMA medium for 3 days at 28 °C. Then its mycelia were removed with a sterile loop and suspended in sterile deionized water and shaken by hand until the clumps were dispersed. The fungal suspension contained 4 × 10⁵ colony-forming units of *Mortierella* sp. per milliliter.

Soil minerals

Samples of four standard minerals (allophane, goethite, kaolinite, and montmorillonite) were provided by Dr. Rollin Jones from the Soil Mineralogy Lab of the University of Hawaii. The sample of allophane was originally collected in Hawaii; the sample of goethite was obtained from Biwabik, Minnesota; the sample of kaolinite was collected from Bath, SC (Ward's Natural Science Establishment Inc., Rochester, NY); and the sample of Montmorillonite was collected from Columbia, MO (Department of Geology, University of Missouri). These minerals were used because they are commonly found in soils and exhibit a wide range of P sorption capacity. For instance, allophane is predominant in volcanic ash soils (Andisols); goethite and kaolinite are commonly found in highly weathered soils (Ultisols and Oxisols); and montmorillonite is predominant in Vertisols and Mollisols.

Preliminary Pi sorption isotherms

The samples of the minerals were passed through a 0.5-mm aperture sieve. Then, Pi sorption isotherms were constructed in triplicates following the procedure developed by Fox and Kamprath (1970). Briefly, aliquots of soil minerals (2 g, dry basis) were transferred into 50-mL plastic centrifuge tubes, and then 20-mL of 0.01 M CaCl₂ solution and graded amounts of dissolved KH₂PO₄ were added. To avoid microbial activity, particularly Pi uptake, the tubes and their contents were autoclaved (120 °C, 0.1 MPa for 30 min). Later, centrifuge tubes and their contents were horizontally shaken (twice a day, 30 min each time) in a reciprocal shaker for 7 days to reach equilibrium of Pi concentration. After this incubation, soil mineral suspension were centrifuged at 2,500 × g for 15 min, and the supernatant was filtered through a Whatman No. 42 filter paper followed by filtration through a Millipore membrane filter (0.45 μm). Solution Pi concentration in the filtrates was determined using the molybdate-blue method (Murphy and Riley 1962).

Microbial Pi desorption experiments

Based on the Pi sorption isotherms, amounts of KH₂PO₄ in 0.01 M CaCl₂ solution were mixed with 2 g of each mineral sample in 50-mL centrifuge tubes to reach the following solution P concentrations: 0.05, 0.1, and 0.2 mg/L (Table 1). Suspension pH was adjusted to 6.0 with 0.1 M NaOH. The tubes were shaken as explained above. Then, the solution was removed by centrifugation at 2,500 × g, and the minerals were washed three times with deionized water. The Pi remaining on the soil minerals was considered as adsorbed. After that, the minerals were oven-dried at 60 °C for 2 days.

At the end of the drying treatment, each tube received 19 mL of a nutrient solution containing 1.0 g NaCl, 0.2 g CaCl₂ 2H₂O, 0.4 g MgSO₄ 7H₂O, 1.0 g NH₄NO₃, and 10.0 g glucose per liter. The sole source of P was the Pi adsorbed on the surfaces of the minerals. Then, the tubes and their contents were autoclaved (120 °C, 30 min, 0.1 MPa). *Mortierella* sp., at the concentration mentioned

Table 1 Amount of P added (in milligrams per liter) to soil minerals to achieve four target solution P concentrations

Initial solution P (mg L ⁻¹)	Soil minerals			
	Montmorillonite	Kaolinite	Goethite	Allophane
0.002–0.006	0 (0.002)	0 (0.003)	0 (0.006)	0 (0.002) ^a
0.05	238	26	143	3,009
0.1	289	51	183	3,578
0.2	340	94	224	4,146

^a Values in parentheses are initial solution P (in milligrams per liter) found in soil minerals and soils without KH₂PO₄ addition

Table 2 Significant *P* values of ANOVA tests for solution Pi concentration as a function of the initial P level and the inoculation with the fungus *Mortierella* sp. in four soil minerals

Source	Montmorillonite	Goethite	Kaolinite	Allophane
Initial P level (A)	<0.0001	<0.0001	<0.0001	<0.0001
Inoculation (B)	<0.0001	<0.0001	<0.0001	NS
A × B	<0.0001	<0.0001	<0.0001	NS
CV (%)	10.7	15.9	11.5	10.7

CV coefficient of variation, NS not significant

above, was aseptically transferred into the tubes at the rate of 1 mL per Erlenmeyer. Tubes not inoculated with the fungus received 19 mL of the nutrient solution and 1 mL of sterilized deionized water. The tubes were continuously shaken on a reciprocal shaker (at 100 cycles per minute) for 6 days at 25 °C.

At the end of the incubation period, the suspensions were centrifuged at 2,500×*g* for 15 min, and the supernatant was filtered through a Whatman no. 42 filter paper followed by filtration through a Millipore membrane filter (0.45 μm). Solution Pi concentration in the filtrates was determined

using the molybdate-blue method (Murphy and Riley 1962). The presence of Pi in the solution was considered an evidence of Pi desorption. The effectiveness of *Mortierella* sp. to desorb Pi was calculated by subtracting Pi desorbed measured in the absence of the fungus from that measured in its presence; this difference was designated as microbially desorbed Pi. The microbial Pi desorption efficiency was calculated based on this difference over the amount of P sorbed. On the other hand, an aliquot of 100 μL from the serial dilution 10⁻² (prepared from the pellets) was aseptically transferred onto a selective YMA medium that contained streptomycin sulfate (500 μg L⁻¹), benomyl (75 μg L⁻¹), and cycloheximide (100 μg L⁻¹). The fungus tolerates both fungicides at these concentrations (Osorio 2008). Then, Petri dishes were incubated at 28 °C for 4 days, and colony-forming units (CFU) were counted.

Experimental design and data analysis

Each mineral was used in a separate experiment. The treatments consisted of a factorial combination (4×2) of four levels of sorbed Pi (corresponding to four solution Pi concentrations) and two levels of inoculation (with or without




Fig. 1 Concentration of P in solution (in milligrams per liter) desorbed from four soil minerals as a function of four initial solution P concentrations and inoculation with *Mortierella* sp. (black

squares, inoculated; white squares, uninoculated). The asterisk means that the two data at a given initial P level are significantly different (LSD test, $P \leq 0.05$)

Mortierella sp.). Treatments were arranged in completely randomized design with three replicates per treatment. Data were subjected to analysis of variance (ANOVA), least significant difference (LSD) tests, regression, and correlation (P value of 0.05) to detect treatment effects. For this purpose, the software Statgraphics Plus version 4.0 (Statpoint, Inc.; Herndon, Virginia) was used.

Results and discussion

In all minerals, except in the allophane, *Mortierella* sp. was capable of increasing solution P concentration significantly (Table 2). However, the effect of the fungus on the concentration of Pi in the solution depended on the amount of Pi initially sorbed on the surface of minerals (Fig. 1a–d).

In the case of allophane, *Mortierella* sp. did not increase solution Pi at any level of sorbed Pi (Fig. 1a). The very high Pi-sorbing capacity of the mineral was apparently responsible for the ineffectiveness of the fungus to desorb Pi. In contrast, *Mortierella* sp. inoculation increased Pi concentration in solution in all other tested soil minerals, except when Pi was not

added to them prior to incubation (Fig 1b–d). The magnitude of Pi desorption by *Mortierella* sp. increased by increasing the amount of sorbed Pi (higher saturation of Pi-sorbing sites).

The effectiveness of *Mortierella* sp. in desorbing Pi from the soil minerals followed the order montmorillonite > kaolinite > goethite > allophane.

The capacity of *Mortierella* sp. to desorb sorbed Pi was demonstrated in three of the four tested minerals (Fig. 1). Generally, the effectiveness of the fungus to desorb Pi depended on the Pi-adsorbing capacity of minerals, which can be expressed as the $P_{0.2}$ value (the amount of P required to increase soluble P concentration up to 0.2 mg L^{-1}). The capacity of *Mortierella* sp. to desorb Pi is correlated ($r=0.87\text{--}0.98$) with the initial level of soluble Pi (as an indicator of the amount of Pi adsorbed on the mineral surface), and it is inversely associated with the Pi sorption capacity of the mineral (Table 3). Therefore, the fungus was ineffective to desorb Pi from the surface of allophane, and probably it may also be ineffective in soils whose mineralogy is dominated by allophane (Andisols).

The capacity of *Mortierella* sp. in releasing Pi from soil minerals can be useful for the fungus as well as other

Table 3 Desorption of P from four soil minerals and microbial P desorption efficiency as a function of initial P concentration, P adsorbed, and P sorption saturation level

Mineral	Initial soluble P (mg L^{-1})	P adsorbed (mg kg^{-1})	P sorption saturation ^a (%)	P desorbed Uninoculated (mg kg^{-1})	P desorbed Inoculated (mg kg^{-1})	Microbial P-desorbed ^b (mg kg^{-1})	Microbial P desorption efficiency ^c (%)
Montmorillonite	0.003	–	–	0.06	0.32	0.26	–
	0.05	238	61.9	0.38	5.1	4.72	1.98
	0.1	289	75.1	0.52	5.9	5.38	1.86
	0.2	340	88.4	0.89	8.8	7.91	2.33
	Microbial P desorbed (mg kg^{-1})= $1.4913+34.7863$ (initial soluble Pi) [$r=0.87$; P value<0.001]						
Kaolinite	0.002	–	–	0.04	0.07	0.03	–
	0.05	26	6.8	0.43	1.40	0.97	3.73
	0.1	51	13.3	0.91	4.17	3.26	6.40
	0.2	94	24.4	1.92	7.04	5.12	5.45
	Microbial P desorbed (mg kg^{-1})= $0.0009+26.641$ (initial soluble Pi) [$r=0.98$; P value<0.001]						
Goethite	0.006	–	–	0.07	0.34	0.27	–
	0.05	143	37.2	0.34	0.67	0.33	0.23
	0.1	183	47.6	0.41	1.78	1.37	0.75
	0.2	224	58.2	0.74	2.53	1.79	0.80
	Microbial P desorbed (mg kg^{-1})= $0.0701+9.2899$ (initial soluble Pi) [$r=0.92$; P value<0.001]						
Allophane	0.002	–	–	0.016	0.022	0.006	–
	0.05	3009	60.2	0.45	0.47	0.02	0.001
	0.1	3578	71.6	0.74	0.92	0.18	0.005
	0.2	4146	82.9	1.40	1.49	0.09	0.002
	Microbial P desorbed was not significant in allophane						

^a P sorption saturation = (P adsorbed/maximal P sorption capacity) \times 100

^b Microbial P desorbed = P desorbed in inoculated treatment – P desorbed in uninoculated treatment

^c Microbial P desorption efficiency = (microbial P desorbed/P adsorbed) \times 100

organisms, including plants and mycorrhizae. This phenomenon may, in part, explain the synergistic interaction of this fungus and the mycorrhizal fungus *Glomus aggregatum* to stimulate Pi uptake and growth of *Leucaena leucocephala* in a highly weathered Oxisol of Hawaii with moderate Pi-sorbing capacity (Osorio and Habte 2001).

The increasing Pi desorption of *Mortierella* sp. by increasing Pi concentration on the mineral may depend on the fact that the strength with which Pi is adsorbed on the mineral surfaces decreases by increasing the saturation of the adsorptive surfaces with Pi (Do Carmo Harta and Torrent 2007). The same probably occurred in several studies conducted with PSM in soils presumably dominated by kaolinite and montmorillonite (Asea et al. 1988; Kucey 1988; Kucey and Leggett 1989; Omar 1998; Peix et al. 2001).

Probably *Mortierella* sp. desorbed Pi by releasing oxalic acid/oxalate (Osorio and Habte 2009; Osorio 2011). It is very well known that organic acids/anions can desorb Pi from sorbing sites (Welch et al. 2002; Jones et al. 2003; Trolove et al. 2003; Ramirez and Osorio 2005; Jara et al. 2006; Sato and Comerford 2006), and their effects in highly weathered soils and in volcanic ash soils containing comparable minerals to those used in the current study have been studied by Hue (1991), Bolan et al. (1994), and Sato and Comerford (2006). The ability of microorganisms to produce organic acids/anions occurs in the rhizosphere where by carbonaceous compounds are released by roots (Amos and Walters 2006; Le Bayon et al. 2006; Marschner et al. 2006; Corrales et al. 2007; Zhang et al. 2011). Once the root exudates are released, they are metabolized by rhizosphere microorganisms leading to the production of organic acids/anions (Gyaneshwar et al. 2002). Although this excretion appears to be genetically determined (Rodriguez et al. 2000, 2006), environmental conditions such as Pi deficiency or high ammonium concentration could trigger or enhance the production of organic acid/anions (Whitelaw 2000; Chen et al. 2006; Vyas et al. 2007).

The efficiency of *Mortierella* sp. to desorb Pi depended on the mineral and the initial P concentration (Table 3). The highest efficiency in desorbing P was detected in kaolinite (3.73–5.45 %), followed by montmorillonite (1.86–2.33 %) and then by goethite (0.23–0.8 %). In general, this efficiency increased with the increase of the amount of P sorbed. This tendency is comparable with that reported by He et al. (2002). On the other hand, in allophane, the P desorption efficiency was extremely low (0.001–0.005 %), which is consistent with the strong P sorption of this soil mineral.

Counts of the fungus were detected only when minerals were inoculated; the lowest counts were found when Pi was not added ($3.0\text{--}4.0 \times 10^3$ CFU g⁻¹ of mineral), likely due to the low available Pi (0.002–0.006 mg L⁻¹) that limited the fungal growth. The two highest level of initial soluble Pi (0.1 and 0.2 mg L⁻¹) produced the highest counts.

This effect was higher in montmorillonite and kaolinite ($8.8\text{--}18.4 \times 10^3$ CFU g⁻¹) than in goethite and allophane ($4.9\text{--}7.6 \times 10^3$ CFU g⁻¹). The number of CFU were correlated with the initial soluble Pi concentration ($r=0.67$, P value=0.004) and, particularly, with the final Pi concentrations ($r=0.96$, $P<0.0001$).

The results presented here indicate the potential effectiveness of *Mortierella* sp. in increasing soil Pi bioavailability by desorbing it from surfaces of soil minerals. We have also established the effectiveness of *Mortierella* sp. in dissolving Pi from rock phosphate (Osorio and Habte 2001, 2009). The effectiveness of this fungus in increasing soil solution Pi by dissolving rock phosphate and desorbing sorbed Pi may play an important role in Pi-deficient soils, particularly in the tropics where the high Pi sorption capacity of soils constrains plant productivity.

Conclusions

This paper showed that (1) the fungus *Mortierella* sp. was effective in desorbing Pi adsorbed on the surface of soil minerals; (2) the effectiveness of the fungus in doing so depended on the Pi sorption capacity on the soil mineral, and this followed the order montmorillonite > kaolinite > goethite > allophane; and (3) the efficiency of the fungus in desorbing Pi increased by increasing the amount of Pi adsorbed on the surface of the mineral.

Acknowledgments We want to thank COLCIENCIAS for the financial support provided to N.W. Osorio during his stay at the University of Hawaii.

References

- Amos B, Walters DT (2006) Maize root biomass and net rhizodeposited carbon: an analysis of the literature. *Soil Sci Soc Am J* 70:1489–1503
- Asea PEA, Kucey RMN, Stewart JWB (1988) Inorganic phosphate solubilisation by two *Penicillium* species in solution culture and soil. *Soil Biol Biochem* 20:459–464
- Bashan Y, Kamnev AA, de-Bashan LE (2012) Tricalcium phosphate is inappropriate as a universal selection factor for isolating and testing phosphate-solubilizing bacteria that enhance plant growth: a proposal for an alternative procedure. *Biol Fertil Soils* (in press) DOI: 10.1007/s00374-012-0737-7
- Bolan NS, Naidu R, Mahimairaja S, Baskaran S (1994) Influence of low-molecular-weight organic acids on the solubilization of phosphates. *Biol Fertil Soils* 18:311–319
- Chen YP, Rekha PD, Arun AB, Shen FT, Lai WA, Young CC (2006) Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Appl Soil Ecol* 34:33–41
- Corrales I, Amenos M, Poschenrieder C, Barcelo J (2007) Phosphorus efficiency and root exudates in two contrasting tropical maize varieties. *J Plant Nutr* 30:887–900

- Do Carmo Harta M, Torrent J (2007) Phosphorus desorption kinetics in relation to phosphorus forms and sorption properties of Portuguese acid soils. *Soil Sci* 172:631–638
- Fox RL, Kamprath E (1970) Phosphate sorption isotherms for evaluating phosphorus requirements of soils. *Soil Sci Soc Am Proc* 34:902–907
- Guppy CN, Menzies NW, Moody PW, Blamey FPC (2005) Competitive sorption reactions between phosphorus and organic matter in soil: a review. *Aust J Soil Res* 43:189–202
- Gyaneshwar P, Kumar GN, Parekh LJ, Poole PS (2002) Role of soil microorganisms in improving P nutrition of plants. *Plant Soil* 245:83–93
- He ZL, Zhu J (1998) Microbial utilization and transformation of phosphate adsorbed by variable charge minerals. *Soil Biol Biochem* 30:917–923
- He ZL, Bian W, Zhu J (2002) Screening and identification of microorganisms capable of utilizing phosphate adsorbed by goethite. *Commun Soil Sci Plant Anal* 33:647–663
- Hoberg E, Marschner P, Lieberei R (2005) Organic acid exudation and pH changes by *Gordonia* sp. and *Pseudomonas fluorescens* grown with P adsorbed to goethite. *Microbiol Res* 160:177–187
- Hue NV (1991) Effects of organic acids/anions on P sorption and phytoavailability in soils with different mineralogies. *Soil Sci* 152:463–471
- Jackman JM, Jones RC, Yost RS, Babcock CJ (1997) Rietveld estimates of mineral percentages to predict phosphate sorption by selected Hawaiian soils. *Soil Sci Soc Am J* 61:618–625
- Jara AA, Violante A, Pigna M, Mora ML (2006) Mutual interactions of sulfate, oxalate, citrate, and phosphate on synthetic and natural allophanes. *Soil Sci Soc Am J* 70:337–346
- Jones DL, Dennis PG, Owen AG, Van Hees PAW (2003) Organic acid behavior in soils—misconceptions and knowledge gaps. *Plant Soil* 248:31–41
- Kirk GJD, Santos EE, Findenegg GR (1999) Phosphate solubilization by organic anion excretion from rice (*Oryza sativa* L.) growing in aerobic soil. *Plant Soil* 211:11–18
- Kucey RMN (1988) Effect of *Penicillium bilaji* on the solubility and uptake of P and micronutrients from soil by wheat. *Can J Soil Sci* 68:261–270
- Kucey RMN, Leggett ME (1989) Microbial mediated increases in plant available phosphorus. *Adv Agron* 42:199–228
- Le Bayon RC, Weisskopf L, Martinoia E, Jansa J, Frossard E, Keller F, Follmi KB, Gobat JM (2006) Soil phosphorus uptake by continuously cropped *Lupinus albus*: a new microcosm design. *Plant Soil* 283:309–321
- Li XF, Zuo FH, Ling GZ, Li YY, Yu YX, Yang PQ, Tang XL (2009) Secretion of citrate from roots in response to aluminum and low phosphorus stresses in *Stylosanthes*. *Plant Soil* 325:219–229
- Marschner P, Solaiman Z, Rengel Z (2006) Rhizosphere properties of Poacea genotypes under P-limiting conditions. *Plant Soil* 283:11–24
- Murphy J, Riley JP (1962) A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27:31–35
- Nwoke OC, Diels J, Abaidoo R, Nziguheba G, Merckx R (2008) Organic acids in the rhizosphere and root characteristics of soybean (*Glycine max*) and cowpea (*Vigna unguiculata*) in relation to phosphorus uptake in poor savanna soils. *Afr J Biotechnol* 7:3620–3627
- Omar SA (1998) The role of rock-phosphate-solubilizing fungi and vesicular-arbuscular-mycorrhiza (VAM) in growth of wheat plants fertilized with rock phosphate. *World J Microb Biot* 14:211–218
- Onweremadu EU (2007) Predicting phosphorus sorption characteristics in highly weathered soils of South-Eastern Nigeria. *Res J Environ Sci* 1:47–55
- Osorio NW (2008) Effectiveness of microbial solubilization of phosphate in enhancing plant phosphate uptake in tropical soils and assessment of the mechanisms of solubilization. Dissertation, University of Hawaii
- Osorio NW (2011) Effectiveness of phosphate solubilizing microorganisms in increasing plant phosphate uptake and growth in tropical soils. In: Maheshwari DK (ed) *Bacteria in agrobiology: plant nutrient management Volume III*. Springer, Berlin, pp 65–80
- Osorio NW, Habte M (2001) Synergistic influence of an arbuscular mycorrhizal fungus and P solubilizing fungus on growth and plant P uptake of *Leucaena leucocephala* in an Oxisol. *Arid Land Res Mgmt* 15:263–274
- Osorio NW, Habte M (2009) Strategies for utilizing arbuscular mycorrhizal fungi and phosphate-solubilizing microorganisms for enhanced phosphate uptake and growth of plants in the soils of the tropics. In: Khan MS, Zaidi A, Musarrat J (eds) *Microbial strategies for crop improvement*. Springer, Berlin, pp 325–351
- Pandey A, Trivedi P, Kumar B, Palni LMS (2006) Characterization of a phosphate solubilizing microorganism and antagonistic strain of *Pseudomonas putida* (B0) isolated from a sub-alpine location in the Indian Central Himalaya. *Curr Microbiol* 53:102–107
- Peix A, Rivas-Boyer AA, Mateos PF, Rodriguez-Barrueco C, Martinez-Molina E, Velasquez E (2001) Growth promotion of chickpea and barley by a phosphate solubilizing strain of *Mesorhizobium mediterraneum* under growth chamber conditions. *Soil Biol Biochem* 33:103–110
- Puente ME, Li CY, Bashan Y (2009) Rock-degrading endophytic bacteria in cacti. *Environ Exp Bot* 66:389–401
- Radersma S, Grierson P (2004) Phosphorus mobilization in agroforestry: organic anions, phosphatase activity and phosphorus fractions in the rhizosphere. *Plant Soil* 259:209–219
- Ramirez F, Osorio W (2005) Effect of low molecular weight organic acids on phosphorus sorption in Andisols. *Suelos Ecuat* 35:71–77 (In Spanish)
- Reddy MS, Kumar S, Babita K, Reddy MS (2002) Biosolubilization of poorly soluble rock phosphates by *Aspergillus tubigenis* and *Aspergillus niger*. *Bioresource Technol* 84:187–189
- Rodriguez H, Fraga R, Gonzalez T, Bashan Y (2006) Genetics of phosphate solubilizing and its potential applications for improving plant growth-promoting bacteria. *Plant Soil* 287:15–21
- Rodriguez H, Gonzalez T, Selman G (2000) Expression of a mineral phosphate solubilizing gene from *Erwinia herbicola* in two rhizobacterial strains. *J Biotechnol* 84:155–161
- Sato S, Comerford N (2006) Organic anions and phosphorus desorption and bioavailability in a humid Brazilian Ultisol. *Soil Sci* 171:695–705
- Shoji S, Nanzyo M, Dahlgren RA (1993) Volcanic ash soils—genesis, properties, and utilization. Elsevier, Amsterdam
- Trolove SN, Hedley MJ, Kirk GJD, Bolan NS, Loganathan P (2003) Progress in selected areas of rhizosphere research on P acquisition. *Aust J Soil Res* 41:471–499
- Vistoso E, Theng BKG, Bolan NS, Parfitt RL, Mora ML (2012) Competitive sorption of molybdate and phosphate in Andisols. *J Soil Sci Plant Nutr* 12:59–72
- Vyas P, Rahi P, Chauhan A, Gulati A (2007) Phosphate solubilization potential and stress tolerance of *Eupenicillium parvum* from tea soil. *Mycol Res* 111:931–938
- Welch S, Taunton AE, Banfield JF (2002) Effect of microorganisms and microbial metabolites on apatite dissolution. *Geomicrobiol J* 19:343–367
- Whitelaw MA (2000) Growth promotion of plants inoculated with phosphate-solubilizing fungi. *Adv Agron* 69:99–151
- Zhang H, Wu X, Li G, Qin P (2011) Interactions between arbuscular mycorrhizal fungi and phosphate-solubilizing fungus (*Mortierella* sp.) and their effects on *Kosteletzkya virginica* growth and enzyme activities of rhizosphere and bulk soils at different salinities. *Biol Fertil Soils* 47:543–554